

Perioperative medicine update

Dr Kate Hudig

Specialist anaesthetist, Auckland City Hospital

We are all used to asking our patients the age-old question “Can you walk up 2 flights of stairs without stopping?” as a way of assessing their cardiorespiratory reserve prior to having surgery. Several single centre studies have raised concerns with the validity of this method for assessing functional capacity^{3,4}. A recent large multicentre cohort study (The METS study) found that subjective physicians’ assessment did not correctly identify those patients who performed poorly on formal pre-operative exercise testing or those at an elevated risk of post-operative morbidity and mortality¹.

Evaluation of functional capacity is considered one of the corner stones of pre-operative anaesthetic risk assessment, guiding both of decisions on further pre-operative testing, optimisation and risk discussion as well as influencing a patient’s intraoperative course and post-operative disposition². With a multi-centre study now confirming concerns that subjective assessment is not an accurate way of assessing this, what should we be doing?

The METS study and several of the METS study sub analyses have helped to answer this question^{1,5,6}. These studies have shown us that using the Duke Activity Status Index (DASI) can help predict post-operative morbidity⁶, while an elevated pre-operative NT-pro BNP is associated with increased mortality at 1 year¹. A sub study looking at the 6 minute walk test, suggested that using a low cut off 370m helped to predict increased morbidity and mortality at 1 year⁶.

The aim of this presentation is to discuss a pragmatic and cost-effective way to assess high-risk patients’ functional capacity before surgery based on current evidence. Can we now put subjective physicians’ assessment to rest?

References:

1. Wijesundera, D.N., Pearse, R.M., Shulman, M.A., Abbott, T.E., Torres, E., Ambosta, A., Croal, B.L., Granton, J.T., Thorpe, K.E., Grocott, M.P. and Farrington, C., 2018. Assessment of functional capacity before major non-cardiac surgery: an international, prospective cohort study. *The Lancet*, 391(10140), pp.2631-2640.
2. Fleisher, L.A., Fleischmann, K.E., Auerbach, A.D., Barnason, S.A., Beckman, J.A., Bozkurt, B., Davila-Roman, V.G., Gerhard-Herman, M.D., Holly, T.A., Kane, G.C. and Marine, J.E., 2014. 2014 ACC/AHA guideline on perioperative cardiovascular evaluation and management of patients undergoing noncardiac surgery: a report of the American College of Cardiology/American Heart Association Task Force on practice guidelines. *Journal of the American College of Cardiology*, 64(22), pp.e77-e137.
3. Reilly, D.F., McNeely, M.J., Doerner, D., Greenberg, D.L., Staiger, T.O., Geist, M.J., Vedovatti, P.A., Coffey, J.E., Mora, M.W., Johnson, T.R. and Guray, E.D., 1999. Self-reported exercise tolerance and the risk of serious perioperative complications. *Archives of internal medicine*, 159(18), pp.2185-2192.
4. Wiklund, R.A., Stein, H.D. and Rosenbaum, S.H., 2001. Activities of daily living and cardiovascular complications following elective, noncardiac surgery. *The Yale journal of biology and medicine*, 74(2), p.75.

5. Shulman, M.A., Cuthbertson, B.H., Wijeyesundera, D.N., Pearse, R.M., Thompson, B., Torres, E., Ambosta, A., Wallace, S., Farrington, C., Myles, P.S. and Ellis, M., 2019. Using the 6-minute walk test to predict disability-free survival after major surgery. *British journal of anaesthesia*, 122(1), pp.111-119.
6. Wijeyesundera, D.N., Beattie, W.S., Hillis, G.S., Abbott, T.E., Shulman, M.A., Ackland, G.L., Mazer, C.D., Myles, P.S., Pearse, R.M., Cuthbertson, B.H. and Myles, P.S., 2020. Integration of the Duke Activity Status Index into preoperative risk evaluation: a multicentre prospective cohort study. *British journal of anaesthesia*, 124(3), pp.261-270.